

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A01N 43/04, 63/00, A61K 31/70, 48/00,</b> <b>C12N 15/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/34202</b> <b>(43) International Publication Date:</b> 21 December 1995 (21.12.95)
<b>(21) International Application Number:</b> PCT/US95/07554 <b>(22) International Filing Date:</b> 14 June 1995 (14.06.95)  <b>(30) Priority Data:</b> 08/260,201      15 June 1994 (15.06.94)      US 08/278,282      21 July 1994 (21.07.94)      US  <b>(71) Applicants:</b> ALEXION PHARMACEUTICALS, INC. [US/US]; Suite 360, 25 Science Park, New Haven, CT 06511 (US). THE AUSTIN RESEARCH INSTITUTE [AU/AU]; Kronheimer Building, Austin Hospital, Studley Road, Heidelberg, VIC 3084 (AU).  <b>(72) Inventors:</b> SANDRIN, Mauro, S.; 211 Barkly Street, Brunswick, VIC 3056 (AU). FODOR, William, L.; 236 Wildcat Road, Madison, CT 06443 (US). ROTHER, Russell, P.; 384 Black's Road, Cheshire, CT 06410 (US). SQUINTO, Stephen, P.; 16 Coachmans Lane, Bethany, CT 06524 (US). MCKENZIE, Ian, F., C.; 359 Brunswick Road, West Brunswick, VIC 3055 (AU).  <b>(74) Agent:</b> KLEE, Maurice, M.; 1951 Burr Street, Fairfield, CT 06430 (US).		<b>(81) Designated States:</b> AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> METHODS FOR REDUCING HYPERACUTE REJECTION OF XENOGRAPTS		
<b>(57) Abstract</b> <p>A method for reducing xenograft rejection is provided wherein a vector directing the expression of a protein having glycosyltransferase activity is introduced into xenogeneic cells. Cells so prepared exhibit a substantial reduction in their binding to naturally occurring preformed human antibodies and are thus protected from hyperacute rejection.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5      METHODS FOR REDUCING HYPERACUTE REJECTION OF XENOGRAFTSCROSS-REFERENCE TO RELATED APPLICATION

10      This application is a continuation-in-part of  
copinging U.S. Patent Application Serial No. 08/260,201  
filed June 15, 1994.

FIELD OF THE INVENTION

15      This invention relates to xenotransplantation. More  
specifically, the invention relates to methods that will  
prevent or reduce hyperacute rejection of xenogeneic  
cells, tissues and organs following transplantation into  
human recipients. The invention provides methods for  
stably reducing the expression on the surface of a  
xenogeneic cell of the non-human antigen known as  
galactose  $\alpha(1,3)$  galactose. This prevents the phenomenon  
20      of antibody-dependent rejection of xenogeneic cells that  
typically follows exposure to human blood, plasma, or  
serum (e.g., following xenotransplantation into a human  
patient) as a result of the binding of preformed natural  
human antibodies to the surfaces of such cells.

25      BACKGROUND OF THE INVENTION

30      Xenotransplantation: Surgical problems related to  
the transplantation of allogeneic organs (i.e., organs  
from donors of the same species as the transplant  
recipient), such as kidney, liver, heart, lung and  
pancreas, have been largely solved, and immunosuppression  
has been improved such that these procedures are now  
routinely performed with a high degree of success (Brent,  
1991). However, a major problem in transplantation  
medicine today is the provision of sufficient allogeneic  
35      donor organs to satisfy the large numbers of patients  
awaiting a transplant. Given the increasing emphasis on  
the costs of dialysis and hospitalization incurred by

patients awaiting transplantation, there is even greater emphasis on the transplantation of donor organs early in the course of disease. Additionally, it is clear that the supply of human donor allografts cannot satisfy this demand. Alternative sources for replacing diseased organs, tissues, or cells, are mechanical devices or animal organs. All clinical transplantations of animal organs have met with failure other than when closely related Old World primate species, such as, baboon, chimp or gorilla, were used as donors. Unfortunately, the supply of potential Old World primate donors is also limited, and ethical considerations further limit the use of organs from such species. Non-primate species, on the other hand, offer a vast potential source of donors.

The most likely donor species for xenotransplantation appears to be the pig (Cooper et al., 1991 and Niekrasz, et al., 1992). This animal is commonly used commercially, and therefore its use will engender fewer ethical problems than the use of primate donors. Furthermore, the pig is considered a highly suitable donor for anatomical and physiological reasons (Cooper et al., 1991 and Niekrasz, et al., 1992).

Immunological Rejection of Xenografts: The rejection of transplanted cells, tissues, or organs may involve both an extremely rapid hyperacute rejection (HAR) phase and a slower cellular rejection phase. HAR of non-human, non-Old World primate organs, tissues, or cells (referred to herein as "xenogeneic" organs, tissues, or cells, or "xenotransplants", or "xenografts") is initiated by preformed natural antibodies found in human blood, plasma, serum, lymph, and the like, that bind to donor cells, e.g., endothelial cells, and activate attack by the complement arm of the human immune system (Dalmaso, et al., 1992; and Tusso, et al., 1993).

While some xenograft tissues (e.g., porcine pancreatic islets) do not appear to be rejected by this mechanism, HAR is the most significant impediment to the

successful xenotransplantation of most cells and tissues, and of all vascularized organs. Methods for the control of the HAR are available. These include interference with the antibody antigen reactions responsible for initiating the HAR response, either by removing the preformed natural antibodies from the circulation or by interference with the binding of the natural antibodies to their specific epitopes (see copending U.S. application Serial No. 08/214,580, entitled "Xenotransplantation Therapies", filed by Mauro S. Sandrin and Ian F.C. McKenzie on March 15, 1994, and PCT publication No. 93/03735, entitled "Methods and Compositions for Attenuating Antibody-Mediated Xenograft Rejection").

A particularly desirable approach to the prevention of hyperacute rejection is to delete or inhibit the  $\alpha(1,3)$  galactosyltransferase gene in xenogeneic cells, and to thus eliminate or significantly reduce expression of Gal  $\alpha(1,3)$  Gal epitopes on the surface of such cells (see copending U.S. patent application Serial No. 08/214,580, supra). This approach eliminates or reduces the binding of preformed natural human antibodies to the xenogeneic cells and, therefore, prevents or reduces the activation of complement and subsequent hyperacute rejection of xenogeneic cells, tissues and organs.

Inhibition of complement attack on the xenotransplant may be accomplished by several means, including the use of complement inhibitors such as the 18kDa C5b-9 inhibitory protein and monoclonal antibodies against human C5b-9 proteins as disclosed in U.S. Patent No. 5,135,916, issued August 4, 1992.

The foregoing methods are effective, but have certain drawbacks in practice, potentially requiring the continuous administration of pharmacologic agents, or, in some cases, requiring the technically difficult production of animals carrying a targeted disruption of a specific gene.

HAR and Complement: Activation of complement leads to the generation of fluid phase (C3a, C5a) and membrane bound (C3b and C5b-9) proteins with chemotactic, procoagulant, proinflammatory, adhesive, and cytolytic properties (Muler-Eberhard, 1988). Immunohistological analysis of hyperacutely rejected xenotransplants reveals antibody deposition, complement fixation, and vascular thrombosis as well as neutrophil infiltration (Zehr, et al., 1994; Auchincloss, 1988; Najarian, 1992; Somerville and d'Apice, 1993; and Mejia-Laguna, et al., 1972).

HAR and Xenoantigens: The targets of natural human antibodies have been the subject of investigations for a number of years, as the identification of these xenoantigens would enable the development of strategies to circumvent hyperacute rejection of xenografts. Several recent studies have convincingly demonstrated that the carbohydrate galactose  $\alpha(1,3)$  galactose (Gal  $\alpha(1,3)$  Gal) is the major xenoepitope recognized by natural human antibodies (see Sandrin, et al., 1993A; Sandrin, et al., 1993B; copending U.S. patent application Serial No. 08/214,580, supra; and PCT publication No. 93/03735, supra).

Galili and colleagues have shown that a large proportion of IgG (1%) in human serum is directed against the Gal  $\alpha(1,3)$  Gal epitope expressed as part of a variety of glycosylated molecules found on both cell surfaces and on secreted glycoproteins (Galili et al., 1984; and Thall and Galili, 1990). This disaccharide epitope is found in all mammals except humans and Old World primates, and naturally occurring preformed anti-Gal  $\alpha(1,3)$  Gal antibodies are found only in humans and Old World primates, i.e., those species which do not themselves express the epitope (Galili et al., 1987 and Galili et al., 1988).

HAR and Preformed Natural Antibodies: The immunoglobulin class of an anti-Gal  $\alpha(1,3)$  Gal antibody determines the biological role of that antibody in



hyperacute rejection. On the basis of histological studies, Bach and Platt (Platt et al., 1990; Platt and Bach 1991; Platt et al., 1991; and Geller et al., 1993) consider that IgM is the most important class of immunoglobulin involved in hyperacute xenograft rejection.

However, natural human antibodies to Gal  $\alpha(1,3)$  Gal are not exclusively of the IgM class, and several studies demonstrate the presence of IgG antibodies reactive with pig cells in human blood (Tusso et al., 1992; Fabian et al., 1992; Hammer et al., 1992; Cairns et al., 1993A; Cairns et al., 1993B; Fournier et al., 1993; Koren et al., 1993; and Zhao et al., 1993), in agreement with the original findings of Galili, et al., 1984 (see also Galili, 1993). For example, by eluting antibodies from different xenogeneic organs after perfusion with normal human serum, Koren et al., 1992, have demonstrated the presence of IgM, IgG and IgA antibodies. Based on these various studies, there is a general consensus that both IgM and IgG antibodies react with Gal  $\alpha(1,3)$  Gal antigens.

The ability of different monosaccharides and oligosaccharides to inhibit the interaction of naturally occurring preformed human antibodies with pig cells and to prevent the antibody-dependent and complement-mediated lysis of pig cells has been examined (Sandrin et al., 1993A; Sandrin et al., 1993B; PCT publication No. 93/03735, supra; and copending U.S. patent application Serial No. 08/214,580, supra).

Inhibition of the binding of such antibodies to xenogeneic cells was obtained with galactose, or with moieties containing terminal galactose in an  $\alpha$  linkage but not a  $\beta$  linkage. Various carbohydrates have also been shown to contain the target epitopes for several types of naturally occurring preformed human antibodies with other specificities (e.g., ABO blood group antibodies). However, no monosaccharide tested, other

than those containing the Gal  $\alpha(1,3)$  Gal epitope, had any inhibitory effect on the binding of naturally occurring preformed human antibodies to xenogeneic cells. Identical inhibition results were obtained when individual human serum samples from blood group A, B, AB or O individuals were used (Sandrin et al., 1993A and Sandrin et al., 1993B).

Similarly, Cooper and colleagues have demonstrated that, of a total of 132 carbohydrates screened for binding to preformed naturally occurring human IgG and IgM antibodies, each of the four carbohydrate molecules that they found could bind such antibodies contained a terminal  $\alpha$  galactose (Good et al., 1992). The four carbohydrates were:

- (1) Gal  $\alpha(1,3)$  Gal  $\beta(1,4)$  GlcNAc,
- (2) Gal  $\alpha(1,3)$  Gal  $\beta(1,4)$  Glc,
- (3) Gal  $\alpha(1,3)$  Gal  $\beta$ , and
- (4) Gal  $\alpha(1,3)$  Gal.

Sugars such as melibiose (a disaccharide containing a terminal galactose in an  $\alpha$  linkage) coupled to a carrier such as SEPHAROSE can be used to purify anti-Gal  $\alpha(1,3)$  Gal antibodies (Galili et al, 1984 and Galili et al., 1985). In some antibody absorption experiments, human serum was passed over the carrier-sugar matrix in order to prepare serum from which the antibodies reactive with the sugar were removed. The results of testing the cytolytic activity of the sera prepared in these experiments indicate that the majority of the cytotoxic antibodies were removed from the serum by these means (Sandrin et al., 1993A; Sandrin et al., 1993B).

In sum, the results of the sugar inhibition studies, the studies of the binding of antibodies to terminal  $\alpha$  galactose-containing molecules, and the studies of the absorption of antibodies by melibiose-SEPHAROSE, all lead to the conclusion that Gal  $\alpha(1,3)$  Gal epitopes are the

most important epitopes detected by naturally occurring human IgG and IgM antibodies.

#### Glycosyltransferases

5 Mammalian cells display a complex variety of carbohydrate antigens on their surfaces. Carbohydrate epitopes are expressed on all mammalian cells by membrane glycoproteins and glycosphingolipids. Profound changes in the structures of these glycoconjugates frequently accompany important biological processes such as  
10 differentiation and development. The types and numbers of carbohydrate epitopes present on cells vary in different species and in different tissues within a given species (Yamakawa and Nagai, 1978).

15 The structures of these carbohydrate moieties are determined largely by the activities of the glycosyltransferases responsible for oligosaccharide synthesis. Therefore, the population of oligosaccharide molecules displayed on the surface of a given mammalian cell is largely determined by the repertoire of  
20 glycosyltransferases active in the cell (Kornfeld and Kornfeld, 1985).


The glycosyltransferases comprise a family of enzymes that transfer sugars from nucleoside diphosphate-sugar conjugates (donor molecules) to acceptor substrate  
25 molecules, forming covalent linkages. Acceptor substrates are often oligosaccharides or oligosaccharide moieties of larger molecules, but may also be specific proteins or lipids. Glycosyltransferases function in a sequential manner, such that the oligosaccharide product  
30 of a transferase activity often becomes the acceptor substrate for subsequent transferase activity. The final result generally contains a linear and/or branched polymer of component monosaccharides linked to one another.

35 Glycosyltransferases differ from each other with respect to the nature of the nucleoside diphosphate-carbohydrate donor, the nature of the acceptor substrate,

and the glycosidic linkage joining the donor sugar to the acceptor substrate (reviewed by Beyer and Hill, 1982). Examples of glycosyltransferases include the following: galactosyltransferases, fucosyltransferases, sialyltransferases, N-acetylglucosaminyltransferases, N-acetylgalactosaminyltransferases, glucosyltransferases, sulfotransferases, acetylases, and mannosyltransferases.

#### Galactosyltransferases

The galactosyltransferases are examples of glycosyltransferases that transfer galactose from a UDP-galactose donor molecule to an acceptor substrate. One such galactosyltransferase, UDP-Gal:Gal  $\beta(1,4)$  Gal NAcGlc  $\alpha(1,3)$  galactosyltransferase (also referred to as  $\alpha(1,3)$  Gal transferase), is a Golgi membrane-bound enzyme that catalyzes the following reaction:



in which R may be a glycoprotein or a glycolipid (Blanken and Van den Eijnden, 1985). The resulting  $\alpha(1,3)$  linked galactose occupies the terminal non-reducing position in N-acetyllactosamine-type carbohydrate chains and, as such, is a non-charged alternative to chain termination by sialic acid. As discussed above, such  $\alpha(1,3)$  Gal structures are the most important epitopes of xenogeneic cells recognized by naturally occurring preformed human antibodies.

$\alpha(1,3)$  Gal transferase and the Gal  $\alpha(1,3)$  Gal  $\beta 1$ -R (herein referred to as Gal  $\alpha(1,3)$  Gal) product of the activity of this enzyme show both species and tissue-specific expression (Galili et al., 1988). The  $\alpha(1,3)$  Gal transferase is widely expressed in a variety of mammalian species, with the notable exception of Old World primates and humans. These mammals do not express the enzyme due to frameshift and nonsense mutations in their genomic sequences encoding this enzyme (Larsen et al., 1990a).

It is believed that humans and Old World primates have high levels of circulating natural preformed antibodies that bind specifically to the Gal  $\alpha(1,3)$  Gal epitope as a consequence of these mutations and the resultant lack of the epitope in humans and Old World primates. The source of antigen exposure responsible for the natural preformed antibodies in these species has not been definitively established, but is believed to be certain bacteria bearing Gal  $\alpha(1,3)$  Gal epitopes that are normally found in the intestines of humans and Old World primates.

The cDNA encoding the pig  $\alpha(1,3)$  galactosyltransferase has been cloned using cross species hybridization (see copending U.S. patent application Serial No. 08/214,580, supra; and Dabkowski et al., 1993). Sequence comparison shows that at the amino acid level there is approximately 75% identity with the murine and approximately 82% identity with the bovine  $\alpha(1,3)$  Gal transferase sequences, with the catalytic domains of the transferases having the highest identity.

#### Fucosyltransferases

The carbohydrate antigens and glycosyltransferases of the human H blood group, as well as the specific details of the biosynthesis and distribution of the H antigen, have been extensively reviewed (see, for example, Lowe, 1991). Several carbohydrates, including those associated with the H antigen, contain the terminal structure Fucose  $\alpha(1,2)$  Galactose. The synthesis of the Fucose  $\alpha(1,2)$  linkage is catalyzed by specific  $\alpha(1,2)$  fucosyltransferase enzymes. The enzymatic activities of these transferases result in the covalent attachment of L-fucose by an  $\alpha(1,2)$  linkage to a variety of acceptor molecules. The H transferase, for example, is a fucosyltransferase that catalyzes a transglycosylation reaction covalently linking a fucose to a specific oligosaccharide acceptor substrate. In this reaction the fucose is derived from the nucleotide sugar donor

5 molecule GDP-fucose and connected by an  $\alpha(1,2)$  linkage to the Galactose residue of Gal  $\beta(1,3)$  GlcNAc-R or Gal  $\beta(1,4)$  GlcNAc-R acceptor substrates (i.e., galactose linked to N-acetylglucosamine in a  $\beta(1,3)$  or a  $\beta(1,4)$  linkage, where R represents a glycoprotein, protein, glycolipid, or lipid).

10 These acceptor substrates are also the acceptor substrates for the  $\alpha(1,3)$  Gal transferase discussed above, although each transferase utilizes a different nucleotide sugar donor molecule (UDP galactose for  $\alpha(1,3)$  Gal transferase vs. GDP fucose for H transferase). The  $\alpha(1,3)$  Gal transferase and the H transferase have now been cloned (see copending U.S. patent application Serial No. 08/214,580, supra; Stanley, 1992; and Lowe, 1991).  
15 The recombinant expression of various glycosyltransferases, including the H transferase, in cells that would be expected to be expressing the  $\alpha(1,3)$  Gal transferase has been reported (see, for example, Lowe, 1991). However, prior to the present invention, the effects of such recombinant expression on the  
20 expression of the Gal  $\alpha(1,3)$  Gal epitope have been unknown.

#### SUMMARY OF THE INVENTION

25 In view of the foregoing, it is an object of this invention to provide genetically modified xenogeneic organs, tissues, and cells that are less prone to hyperacute rejection when exposed to human blood, plasma, serum, lymph, or the like (e.g., following xenotransplantation into human patients) than their  
30 unmodified precursors, and to provide methods for the preparation of such xenogeneic organs, tissues, and cells. In accordance with these methods, xenogeneic cells are genetically modified so that they express the glycosyltransferase activity of an exogenous  
35 glycosyltransferase (i.e., a glycosyltransferase encoded by a recombinant nucleic acid molecule introduced into

the xenogeneic cells or a parent cell of the xenogeneic cells). In particular, the genetically-modified xenogeneic cells of the invention exhibit reduced levels of the xenoantigen Gal  $\alpha(1,3)$  Gal on their cell surfaces.

5 In another aspect of the invention, the genetically-modified xenogeneic cells of the invention are inhibited from binding to preformed naturally occurring human antibodies and are therefore significantly less sensitive to HAR as demonstrated by reduced sensitivity to  
10 activation and/or lysis by human complement. In this way, when transplanted into human patients, the rejection of such cells by complement-mediated hyperacute rejection mechanisms is reduced or prevented.

15 In certain preferred embodiments, the invention provides a method for reducing rejection of a xenogeneic cell following transplantation into a human or an Old World primate comprising:

(a) producing a genetically altered cell by introducing an expression vector comprising a nucleic  
20 acid sequence encoding a protein having fucosyltransferase activity into a recipient cell, the introduction of said expression vector causing a substantial reduction in the binding of naturally occurring preformed human antibodies or naturally  
25 occurring preformed Old World primate antibodies to the genetically altered cell when compared to the binding of said antibodies to the recipient cell; and

(b) transplanting said genetically altered cell or a cell derived from said cell into a human or an Old  
30 World primate.

In other preferred embodiments the invention provides an ungulate cell which has been genetically altered by the introduction of an expression vector comprising a nucleic acid sequence encoding a protein  
35 having fucosyltransferase activity into a recipient ungulate cell, the introduction of said expression vector causing a substantial reduction in the binding of

5 naturally occurring preformed human antibodies or naturally occurring preformed Old World primate antibodies to said genetically altered ungulate cell when compared to the binding of said antibodies to the recipient ungulate cell.

10 In further preferred embodiments, the invention provides a retroviral packaging or producer cell which has been genetically altered by the introduction of an expression vector comprising a nucleic acid sequence encoding a protein having fucosyltransferase activity into a recipient cell from which the genetically altered retroviral packaging or producer cell is derived, the introduction of said expression vector causing a substantial reduction in the binding of naturally occurring preformed human antibodies or naturally occurring preformed Old World primate antibodies to said genetically altered retroviral packaging or producer cell when compared to the binding of said antibodies to the recipient cell from which the genetically altered retroviral packaging or producer cell is derived.

#### BRIEF DESCRIPTION OF THE DRAWINGS

25 FIGS. 1-4 are photomicrographs of African Green Monkey COS cells which have been fluorescently stained with anti-H antigen mAbs (FIGS. 2 and 3) or with lectins specific for the Gal  $\alpha(1,3)$  Gal epitope (FIGS. 1 and 4). In each figure, the bottom panel shows all cells, as seen by phase contrast illumination, and the top panel shows only those cells specifically binding to the mAb or lectin as seen by ultraviolet illumination. The cells in 30 FIG. 1 have been transfected with a vector expressing the Gal  $\alpha(1,3)$  Gal transferase; the cells in FIG. 2 have been transfected with a vector expressing H transferase; and the cells in FIGS. 3 and 4 have been transfected with equal amounts of both vectors. African Green Monkeys are 35 Old World primates and thus their cells, including COS cells, do not express the Gal  $\alpha(1,3)$  Gal epitope. In addition, COS cells do not express the H epitope.



FIG. 5 illustrates the expression of the H epitope and reduced expression of the Gal  $\alpha(1,3)$  Gal epitope in stably transfected porcine kidney cells as analyzed by lectin staining and fluorescence-based flow cytometric analysis of the cells.

FIG. 6 demonstrates the loss of human serum IgG and IgM binding to porcine kidney cells associated with the expression of the H epitope and reduced expression of the Gal  $\alpha(1,3)$  Gal epitope as analyzed by fluorescence staining and flow cytometry.

FIG. 7 illustrates the enhanced resistance to human serum lysis associated with the expression of the H epitope and consequent reduced expression of the Gal  $\alpha(1,3)$  Gal epitope in stably transfected porcine kidney cells.

The foregoing drawings, which are incorporated in and constitute part of the specification, illustrate the preferred embodiments of the invention, and together with the description, serve to explain the principles of the invention. It is to be understood, of course, that both the drawings and the description are explanatory only and are not restrictive of the invention.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

Glycosyltransferases: A variety of nucleic acid molecules encoding glycosyltransferases can be used in the practice of the present invention provided that the glycosyltransferase encoded by the nucleic acid molecule is able to reduce the levels of Gal  $\alpha(1,3)$  Gal epitopes on the surface of a xenogeneic cell in which the exogenous transferase is expressed. In accordance with the present invention, this property can be determined by introducing an appropriate expression vector directing the expression of the candidate glycosyltransferase into xenogeneic cells and then testing the cells for cell surface levels of the Gal  $\alpha(1,3)$  Gal epitope using, for example, human serum, as described below in Example 4.

Transferases suitable for use in the methods and cells of the present invention will cause a substantial reduction in the binding of naturally occurring preformed human antibodies to the xenogeneic cells after introduction of the expression vector compared to binding before introduction of the vector. An at least 50% reduction in binding will, in general, comprise a "substantial reduction". Smaller reductions in binding are also considered "substantial" if they represent a statistically significant reduction, i.e., a reduction that, when analyzed by a standard statistical test, such as the student's T test, will give a probability value,  $p$ , less than or equal to 0.05 and, preferably, less than or equal to 0.015. Examples of the construction of such vectors, production of such cells, and the testing of such cells for reduction of preformed natural antibody binding are given below in Examples 1-5.

In particular, reduction in the binding of naturally occurring preformed antibodies can be determined by staining and counting stained cells as described below in Example 2, or by FACS analysis as described in Example 4 below in which case a quantitative readout can be obtained by measuring the areas under the various FACS curves and the shifts in the positions of those curves, or by measurement of changes in complement resistance as described in Example 5 below.

While not wishing to be bound by any particular theory of operation, it is believed that glycosyltransferases, which are able to reduce the levels of Gal  $\alpha(1,3)$  Gal epitopes on the surface of a xenogeneic cell in which the transferase is expressed, effect this reduction by competition for a shared acceptor substrate. Specifically, it is believed that transferases suitable for use in the methods and cells of the invention transfer donor sugars to  $\beta(1,3)$  Glc NAc-R or  $\beta(1,4)$  Glc NAc-R acceptor substrates. Thus, preferred transferases include those that transfer donor sugars to  $\beta(1,3)$  Glc

NAC-R or  $\beta(1,4)$  Glc NAC-R acceptor substrates and create covalent linkages other than the Gal  $\alpha(1,3)$  Gal linkage upon such transfer.

Preferred transferases to be used in the practice of the invention include fucosyltransferases. With regard to these transferases, it is believed that the addition of a terminal fucose residue to the  $\beta(1,3)$  Glc NAC-R or  $\beta(1,4)$  Glc NAC-R acceptor substrate of the  $\alpha(1,3)$  galactosyltransferase prevents the addition of a Gal  $\alpha(1,3)$  Gal epitope to the acceptor substrate. Specific examples of fucosyltransferases that can be tested for use in the process of the present invention include the  $\alpha(1,2)$  fucosyltransferase (H transferase; Larsen et al., 1990A) and the  $\alpha(1,3/1,4)$  fucosyltransferase (Weston et al., 1992). Of these, the H transferase is preferred and the human H transferase is particularly preferred. This transferase is responsible for synthesis of the H antigen which is the universal donor O-blood group antigen and utilizes the same acceptor substrates as the  $\alpha(1,3)$  Gal transferase. Alternatively, although less preferred, sialyltransferases, e.g., the  $\alpha(2,6)$  sialyltransferase (see Lowe, 1991), may be used in the practice of the invention.

Although the foregoing discussion and those that follow are phrased in terms of nucleic acid molecules encoding glycosyltransferases, the cells and methods of the invention more generally comprise nucleic acid molecules encoding any and all proteins that have glycosyltransferase activity, including, in particular, fucosyltransferase activity. Such proteins may be in the form of intact glycosyltransferases, but may also be in the form of proteins comprising active mutant glycosyltransferases such as those comprising active fragments of glycosyltransferases. See, for example, Kukowska, et al., 1991.

Vectors for expression of recombinant glycosyltransferases: In addition to the foregoing, the present invention provides vectors for the expression of recombinant glycosyltransferases in xenogeneic cells at levels effective to reduce the expression of Gal  $\alpha(1,3)$  Gal epitopes by the xenogeneic cells into which the vectors have been introduced. Recombinant polynucleotides encoding glycosyltransferases that are appropriate for use in such vectors include those encoding the transferases discussed above. A particularly preferred polynucleotide is that encoding human H transferase, SEQ ID NO: 3.

The nucleic acid encoding the desired exogenous glycosyltransferase may be inserted into an appropriate parent expression vector, i.e., an expression vector that contains a site for inserting protein-encoding nucleic acid molecules, and also contains (in the appropriate orientation for expression) the necessary elements for the transcription and translation of an inserted protein-encoding sequence. Particularly preferred transcriptional and translational signals allow for expression of the desired glycosyltransferase in a wide variety of xenogeneic cell types.

A candidate parent expression vector can be tested for suitability for use in the practice of the present invention by the insertion of a nucleic acid fragment encoding the human H transferase into a site appropriate for expression in the parent expression vector, as described below in Example 1 for the APEX-1 vector, and testing cells containing the resulting expression vector for susceptibility to human complement-mediated damage as described below in Example 5.

The transcriptional and translational control sequences in mammalian expression vector systems to be used in genetically altering vertebrate cells may be provided by various sources, including viral sources. For example, commonly used promoters and enhancers known

to be generally operable in many mammalian cell types are derived from Polyoma virus, Adenovirus, Simian Virus 40 (SV40), and human cytomegalovirus immediate-early gene 1 promoter and enhancer (CMV). Cell-type specific promoters may also be used to express glycosyltransferases in particular cell types if desired.

A particularly preferred eukaryotic vector for the expression of glycosyltransferases in the methods and cells of the invention is pAPEX-1, SEQ ID NO:4 (see also copending U.S. patent application Serial No. 08/252,493, filed June 1, 1994, entitled "Porcine E-Selectin"). pAPEX-1 is a derivative of the vector pcDNA1/Amp (Invitrogen, San Diego CA) which was modified to increase protein expression in mammalian cells (see Example 1, below).

Whatever parent expression vector is used, transferase-encoding polynucleotide fragments will be subcloned into the parent vector, typically following digestion with appropriate restriction endonucleases. Fragments for such subcloning can be obtained by PCR amplification, restriction endonuclease digestion, and the like. These fragments and the parent vectors are assembled into a transferase expression vector using standard methods such as PCR fusion or enzymatic ligation (Sambrook, et al., 1989; Ausubel et al., 1992). Alternatively, nucleic acid molecules encoding the glycosyltransferases used in the methods and cells of the invention can be synthesized by chemical means (Talib, et al., 1991).

Expression vectors preferably also contain selectable markers, such as a beta lactamase antibiotic resistance gene for plasmid selection and propagation in microbial cells in the presence of an antibiotic such as ampicillin, and the neomycin gene for selection and propagation of stable mammalian transfectants, e.g., in the presence of the cytotoxic aminoglycoside G418.

Introduction of nucleic acid molecules into cells via transfection or transduction: As known in the art, introduction of nucleic acid molecules into cells can be accomplished by numerous methods, typically by transfection or transduction. Transfection methods include the addition of chemical carriers such as DEAE/dextran, calcium phosphate, or amphipathic lipids (in which case the procedure is generally referred to in the art as lipofection) to the nucleic acid molecules before or during the addition of those molecules to the cells to be transfected. Transfection methods also include mechanical means, such as electroporation, electric field mediated transfer (also referred to as Baekonization, see, for example, U.S. Pat. No. 4,849,355, entitled "Method Of Transferring Genes Into Cells" and U.S. Pat. No. 4,663,292, entitled "High-Voltage Biological Macromolecular Transfer And Cell Fusion System"), microinjection, and ballistic particle, "gene gun", mediated transfer.

The introduction of nucleic acid molecules into cells using engineered, replication incompetent viruses is referred to in the art as transduction. Transduction is a preferred method of nucleic acid molecule introduction into xenogeneic endothelial cells. The first step needed to use transduction methods in the practice of the present invention is incorporating the genetic sequence of the glycosyltransferase into a viral vector, e.g., a retroviral vector. Thereafter, the retroviral vectors are incorporated into retroviral vector particles using packaging cells.

The manipulation of retroviral nucleic acids to construct retroviral vectors and packaging cells is accomplished using techniques known in the art. See, for example, Ausubel, et al., 1992, Volume 1, Section III (units 9.10.1 - 9.14.3); Sambrook, et al., 1989; Miller, et al., 1989; Eglitis, et al., 1988; U.S. Patents Nos. 4,650,764, 4,861,719, 4,980,289, 5,122,767, and

5,124,263; as well as PCT Patent Publications Nos. WO 85/05629, WO 89/07150, WO 90/02797, WO 90/02806, WO 90/13641, WO 92/05266, WO 92/07943, WO 92/14829, and WO 93/14188.

5 In particular, retroviral vectors for use in the practice of the invention can be prepared and used as follows. First, a retroviral vector comprising a nucleic acid sequence encoding a glycosyltransferase is constructed from a parent retroviral vector. Examples of  
10 such parent retroviral vectors are found in, for example, Korman, et al., 1987; Morgenstern, et al., 1990; U.S. Patents Nos. 4,405,712, 4,980,289, and 5,112,767; and PCT Patent Publications Nos. WO 85/05629, WO 90/02797, and WO 92/07943. A preferred parent retroviral vector is the  
15 Moloney murine leukemia virus-derived expression vector pLXSN (Miller, et al., 1989).

The parent retroviral vector used in the practice of the present invention will be modified to include a glycosyltransferase encoding sequence and will be  
20 packaged into non-infectious (replication incompetent) transducing retroviral particles (virions) using an amphotropic packaging system, preferably one suitable for use in gene therapy applications.

Examples of useful packaging systems are found in,  
25 for example, Miller, et al., 1986; Markowitz, et al., 1988; Cosset, et al., 1990; U.S. Patents Nos. 4,650,764, 4,861,719, 4,980,289, 5,122,767, and 5,124,263, and PCT Patent Publications Nos. WO 85/05629, WO 89/07150, WO 90/02797, WO 90/02806, WO 90/13641, WO 92/05266, WO  
30 92/07943, WO 92/14829, and WO 93/14188. A preferred packaging cell is the PA317 packaging cell line (ATCC CRL 9078).

The generation of "producer cells" is accomplished by introducing retroviral vectors into the packaging  
35 cells. The producer cells generated by the foregoing procedures are used to produce the retroviral vector

particles. This is accomplished by culturing of the cells in a suitable growth medium.

5 Preferably, the virions are harvested from the culture and administered to the target cells which are to be transduced. Examples of such target cells include isolated xenogeneic cells, cells of a xenogeneic organ or tissue, and other cells to be protected from antibody binding and complement attack, as well as xenogeneic progenitor cells, including stem cells such as embryonic or hematopoietic stem cells, which can be used to generate transgenic cells, tissues, or organs.

10 Alternatively, when practicable, virions are added to the target xenogeneic cells to be transduced by co-culture of the target cells with the producer cells. Suitable buffers and conditions for stable storage and subsequent use of the virions can be found in, for example, Ausubel, et al., 1992.

15 Cells, tissues, and organs: In general, any xenogeneic cell, tissue or organ may be utilized in the practice of the present invention. Preferred cells are of ungulate origin, and particularly preferred cells are of pig origin. The glycosyltransferase nucleic acid constructs of the invention can be used to engineer cultured cells of various types for subsequent use in transplantation. Examples of useful cell types include endothelial cells, fibroblastic and other skin cells, hepatic cells, neuronal and glial cells, pancreatic islet cells, hematopoietic cells, blood cells, lens cells, corneal cells, and stem cells.

20 Further, the glycosyltransferase nucleic acid constructs of the invention can be used to alter retroviral packaging cells or retroviral producer cells so that such cells exhibit a substantial reduction in the binding of naturally occurring preformed human antibodies or naturally occurring preformed Old World primate antibodies when compared to the binding of said antibodies to packaging or producer cells which have not

25

30

35



been so altered. In the discussion that follows, the expression "altered retroviral packaging/producer cells" is used to describe either or both of said altered packaging or producer cells. Such altered retroviral packaging/producer cells may be from any species that expresses Gal  $\alpha(1,3)$  Gal epitopes, including cells of rodent or canine origin.

Among other applications, such altered retroviral packaging/producer cells may be used to provide gene therapy treatment in a patient in need of such treatment, e.g., for therapeutic control of neoplastic tumors. In this embodiment of the invention, altered retroviral producer cells producing a retroviral vector particle providing a therapeutic benefit are implanted into the patient. In the case of cancer therapy the implantation is preferably made into or adjacent to the tumor. In accordance with the invention, such altered producer cells are protected from HAR upon transplantation (implantation) into a human or Old World primate patient.

In addition, as disclosed in copending U.S. patent application Serial No. 08/278,639, entitled "Retroviral Transduction of Cells in the Presence of Complement", which is being filed concurrently herewith in the names of Russell P. Rother, Scott A. Rollins, William L. Fodor, and Stephen P. Squinto, the retroviral particles produced by the altered producer cells are protected from inactivation by complement in the body fluids of the patient. Other methods to protect retroviral vector particles from inactivation by complement in the body fluids of humans or Old World primates include those discussed in copending U.S. patent application Serial No. 08/278,550, entitled "Retroviral Transduction of Cells Using Soluble Complement Inhibitors", which is being filed concurrently herewith in the names of Russell P. Rother, Scott A. Rollins, James M. Mason, and Stephen P. Squinto, and in copending U.S. patent application Serial No. 08/278,630, entitled "Retroviral Vector Particles

Expressing Complement Inhibitor Activity", which is also being filed concurrently herewith in the names of James M. Mason and Stephen P. Squinto.

5 General discussions of packaging cells, retroviral vector particles and gene transfer using such particles can be found in various publications including PCT Patent Publication No. WO 92/07943, EPO Patent Publication No. 178,220, U.S. Patent No. 4,405,712, Gilboa, 1986; Mann, et al., 1983; Cone and Mulligan, 1984; Eglitis, et al., 10 1988; Miller, et al., 1989; Morgenstern and Land, 1990; Eglitis, 1991; Miller, 1992; Mulligan, 1993, and Ausubel, et al., 1992. The manipulation of retroviral nucleic acids to construct packaging vectors and packaging cells is discussed in, for example, Ausubel, et al., Volume 1, 15 Section III (units 9.10.1 - 9.14.3), 1992; Sambrook, et al., 1989; Miller, et al., 1989; Eglitis, et al., 1988; U.S. Patents Nos. 4,650,764, 4,861,719, 4,980,289, 5,122,767, and 5,124,263; as well as PCT Patent Publications Nos. WO 85/05629, WO 89/07150, WO 90/02797, 20 WO 90/02806, WO 90/13641, WO 92/05266, WO 92/07943, WO 92/14829, and WO 93/14188. To form packaging cells, packaging vectors are introduced into suitable host cells such as those found in, for example, Miller and Buttimore, Mol. Cell Biol., 6:2895-2902, 1986; Markowitz, et al., J. Virol., 62:1120-1124, 1988; Cosset, et al., J. Virol., 64:1070-1078, 1990; U.S. Patents Nos. 4,650,764, 4,861,719, 4,980,289, 5,122,767, and 5,124,263, and PCT Patent Publications Nos. WO 85/05629, WO 89/07150, WO 90/02797, WO 90/02806, WO 90/13641, WO 92/05266, WO 30 WO 92/07943, WO 92/14829, and WO 93/14188. Once a packaging cell line has been established, producer cells are generated by introducing retroviral vectors into the packaging cells. Examples of such retroviral vectors are found in, for example, Korman, et al., 1987, Proc. Natl. Acad. Sci. USA, 84:2150-2154; Miller and Rosman, 35 Biotechniques, 7:980-990, 1989; Morgenstern and Land, 1990; U.S. Patents Nos. 4,405,712, 4,980,289, and

5,112,767; and PCT Patent Publications Nos. WO 85/05629, WO 90/02797, and WO 92/07943. The retroviral vector includes a psi site and one or more exogenous nucleic acid sequences selected to perform a desired function, e.g., an experimental, diagnostic, or therapeutic function. These exogenous nucleic acid sequences are flanked by LTR sequences which function to direct high efficiency integration of the sequences into the genome of the ultimate target cell. (See also the discussion of transduction set forth above.)

Many applications of gene therapy using retroviral vector particles (RVVPs) are known and have been extensively reviewed (see, for example, Boggs, 1990; Kohn, et al., 1989; Lehn, 1990, Verma, 1990; Weatherall, 1991; and Felgner and Rhodes, 1991).

A variety of genes and DNA fragments can be incorporated into RVVPs for use in gene therapy. These DNA fragments and genes may encode RNA and/or protein molecules which render them useful as therapeutic agents. Protein encoding genes of use in gene therapy include those encoding various hormones, growth factors, enzymes, lymphokines, cytokines, receptors, and the like.

Among the genes which can be transferred are those encoding polypeptides that are absent, are produced in diminished quantities, or are produced in mutant form in individuals suffering from a genetic disease. Other genes of interest are those that encode proteins that, when expressed by a cell, can adapt the cell to grow under conditions where the unmodified cell would be unable to survive, or would become infected by a pathogen. Genes encoding proteins that have been engineered to circumvent a metabolic defect are also suitable for transfer into the cells of a patient. Such genes include the transmembrane form of CD59 discussed in copending U.S. patent application No. 08/205,720, filed March 3, 1994, entitled "Terminal Complement Inhibitor Fusion Genes and Proteins" and copending U.S. patent

application No. 08/206,189, filed March 3, 1994, entitled "Method for the Treatment of Paroxysmal Nocturnal Hemoglobinuria".

5 In addition to protein-encoding genes, RVVPs can be used to introduce nucleic acid sequences encoding medically useful RNA molecules into cells. Examples of such RNA molecules include anti-sense molecules and catalytic molecules, such as ribozymes.

10 In order to expedite rapid transduction by eliminating the need to wait for target cells to divide, and to allow transduction of cells that divide slowly or not at all, the use of RVVPs that can transduce non-dividing cells may be preferred. Such RVVPs are disclosed in copending U.S. patent applications Serial  
15 Nos. 08/181,335 and 08/182,612, both entitled "Retroviral Vector Particles for Transducing Non-Proliferating Cells" and both filed January 14, 1994. These patent applications also discuss specific procedures suitable for producing packaging vectors and retroviral vectors as  
20 well as the use of such vectors to produce packaging cells and producer cells, respectively.

Transgenic animals: Transgenic xenogeneic animals provide a preferred source of the cells, tissues, and organs of the invention. In accordance with certain  
25 aspects of the invention, the nucleic acid molecules of the invention are used to generate engineered transgenic animals, preferably ungulates (i.e., hooved animals such as pigs, cows, goats, sheep, and the like), that express the carbohydrate products of glycosyltransferases on the  
30 surfaces of their cells (e.g., endothelial cells) using techniques known in the art.

These techniques include, but are not limited to, microinjection (e.g., of pronuclei), electroporation of ova or zygotes, electric field mediated transfer (i.e.,  
35 Baekonization, supra; see also Zhao and Wong, 1991), nuclear transplantation, and/or the stable transfection or transduction of embryonic stem cells derived from the

animal of choice. Electric field mediated transfer, i.e., Baekonization, is a preferred method of producing the transgenic animals of the invention.

5 A common element of these techniques involves the preparation of a transgene transcription unit. Such a unit comprises a DNA molecule which generally includes: 1) a promoter, 2) the nucleic acid sequence of interest, i.e., the sequence encoding a glycosyltransferase, and 3) a polyadenylation signal sequence. Other sequences, such  
10 as enhancer and intron sequences, can be included if desired. The unit can be conveniently prepared by isolating a restriction fragment of a plasmid vector which expresses the glycosyltransferase protein in, for example, mammalian cells. Preferably, the restriction  
15 fragment is free of sequences which direct replication in bacterial host cells since such sequences are known to have deleterious effects on embryo viability.

The most well known method for making transgenic animals is that used to produce transgenic mice by  
20 superovulation of a donor female, surgical removal of the egg, injection of the transgene transcription unit into the pro-nuclei of the embryo, and introduction of the transgenic embryo into the reproductive tract of a pseudopregnant host mother, usually of the same species.  
25 See Wagner, U.S. Patent No. 4,873,191, Brinster, et al., 1985, Hogan, et al., 1986, Robertson 1987, Pedersen, et al., 1990.

The use of this method to make transgenic ungulates is also widely practiced by those of skill in the art.  
30 As an example, transgenic swine are routinely produced by the microinjection of a transgene transcription unit into pig embryos. See, for example, PCT Publication No. WO92/11757. In brief, this procedure may, for example, be performed as follows.

35 First, the transgene transcription unit is gel isolated and extensively purified through, for example, an ELUTIP column (Schleicher & Schuell, Keene, NH),

dialyzed against pyrogen free injection buffer (10mM Tris, pH7.4 + 0.1mM EDTA in pyrogen free water) and used for embryo injection.

5       Embryos are recovered from the oviduct of a  
hormonally synchronized, ovulation induced sow,  
preferably at the pronuclear stage. They are placed into  
a 1.5 ml microfuge tube containing approximately 0.5 ml  
of embryo transfer media (phosphate buffered saline with  
10% fetal calf serum). These are centrifuged for 12  
10   minutes at 16,000 x g in a microcentrifuge. Embryos are  
removed from the microfuge tube with a drawn and polished  
Pasteur pipette and placed into a 35 mm petri dish for  
examination. If the cytoplasm is still opaque with lipid  
such that the pronuclei are not clearly visible, the  
15   embryos are centrifuged again for an additional 15  
minutes.

Embryos to be microinjected are placed into a drop  
of media (approximately 100  $\mu$ l) in the center of the lid  
of a 100 mm petri dish. Silicone oil is used to cover  
20   this drop and to fill the lid to prevent the medium from  
evaporating. The petri dish lid containing the embryos  
is set onto an inverted microscope equipped with both a  
heated stage (37.5-38°C) and Hoffman modulation contrast  
optics (200X final magnification).

25       A finely drawn and polished micropipette is used to  
stabilize the embryos while about 1-2 picoliters of  
injection buffer containing approximately 200-500 copies  
of the purified transgene transcription unit is delivered  
into the nucleus, preferably the male pronucleus, with  
30   another finely drawn and polished micropipette. Embryos  
surviving the microinjection process as judged by  
morphological observation are loaded into a polypropylene  
tube (2 mm ID) for transfer into the recipient  
pseudopregnant sow.

35       Offspring are tested for the presence of the  
transgene by isolating genomic DNA from tissue removed  
from the tail of each piglet and subjecting this genomic

DNA to nucleic acid hybridization analysis with transgene specific probes or PCR analysis with transgene specific primers.

5 Another commonly used technique for generating transgenic animals involves the genetic manipulation of embryonic stem cells (ES cells) as described in PCT Patent Publication No. WO 93/02188 and Robertson, 1987.

10 In accordance with this technique, ES cells are grown as described in, for example, Robertson, 1987, and in U.S. Patent No. 5,166,065 to Williams et al., 1988. Genetic material is introduced into the embryonic stem cells by, for example, electroporation according, for example, to the method of McMahon, et al., 1990, or by transduction with a retroviral vector according, for example, to the  
15 method of Robertson, et al., 1986, or by any of the various techniques described by Lovell-Badge, 1987.

Chimeric animals are generated as described, for example, in Bradley, 1987. Briefly, genetically modified ES cells are introduced into blastocysts and the modified  
20 blastocysts are then implanted in pseudo-pregnant female animals. Chimeras are selected from the offspring, for example by the observation of mosaic coat coloration resulting from differences in the strain used to prepare the ES cells and the strain used to prepare the blastocysts, and are bred to produce non-chimeric  
25 transgenic animals.

Other methods for the production of transgenic animals are disclosed in U.S. Patent No. 5,032,407 to Wagner et al., and PCT Publication No. WO90/08832.

30 Among other applications, transgenic animals prepared in accordance with the invention are useful as model systems for testing the xenotransplantation of their engineered cells, tissues, and organs and as sources of engineered cells, tissues, and organs for  
35 xenotransplantation. The expression of functional glycosyltransferases by endothelial cells and/or other cell types in the tissues and organs of the transgenic

animals of the present invention will provide reduced susceptibility to hyperacute complement-mediated rejection following exposure of those cells, tissues, and organs to complement in human blood, plasma, serum, lymph, or the like, e.g., following xenotransplantation into humans or Old World primates. In accordance with the invention, reduced susceptibility to HAR is provided because naturally occurring preformed human or Old World primate antibodies have fewer binding sites on the transgenic cells of the invention.

Without intending to limit it in any manner, the present invention will be more fully described by the following examples.

#### Example 1

##### H Transferase

The human H transferase gene was cloned from cDNA prepared from Human Epidermoid Carcinoma cells (HEC cells, ATCC CRL 1555 #A-431) utilizing the Polymerase Chain Reaction (PCR). Cytoplasmic RNA was prepared from approximately  $5 \times 10^6$  cells, and first strand cDNA was synthesized from 5 $\mu$ g of RNA in a final volume of 100 $\mu$ l using the following reaction conditions: 10mM Tris-HCl pH8.3; 50mM KCl; 1.5mM MgCl<sub>2</sub>; 500ng oligo(dT)<sub>15</sub> (Promega Corporation, Madison, Wisconsin); 10mM DTT; 0.25mM dNTPs (dG, dC, dA, dT); and 20U Avian Myeloblastosis Virus reverse transcriptase (Seikagaku of America, Inc., Rockville, Maryland) at 42°C for one hour.

PCR was performed following cDNA synthesis using 4 $\mu$ l of first strand cDNA reaction mixture as template and the following primers: a 34 base 5' primer homologous to the 5' untranslated region of the H transferase cDNA (SEQ ID NO:1; 5'-GGCCACGAAA AGCGGACTGT GGATCCGCCA CCTG-3'), where the underlined sequence represents a unique BamHI site; and a 38 base 3' primer homologous to the 3' UTR of the H transferase cDNA (SEQ ID NO:2; 5'-CAGGAACACC ACCAAGCTTC TCGAGAAGATGC CAGGCC-3'), in which the underlined sequence represents a unique XhoI site. PCR



reactions consisted of 35 cycles of 95°C - 1 minute, 52°C - 1 minute, and 72°C - 1.5 minutes. These 35 cycles were followed by a single ten minute extension at 72° C. An approximately 1300 bp band representing the PCR product was seen following agarose gel electrophoresis of an aliquot of the PCR reaction. This PCR product was cloned into a plasmid vector using the T/A cloning kit (Invitrogen, San Diego, CA). The pCRII plasmid vector included in this kit served as the recipient, and the resulting plasmid construct was amplified in E. coli and purified. Positive clones were identified by restriction endonuclease digestion and the insert was subsequently sequenced to confirm that the plasmid construct contained the human H transferase cDNA sequence shown in SEQ ID NO:3. An approximately 1200 bp BamHI-XhoI DNA fragment, encoding the full length H transferase enzyme, was gel isolated from the pCRII plasmid construct, electroeluted and subcloned into a BamHI-XhoI cut pAPEX-1 expression vector (see the following paragraph for a detailed description of this vector). Positive clones were identified by restriction mapping with BamHI-XhoI and StuI. Plasmid pAPEX1-HT, referred to hereinafter as pHT, was the result of these cloning and subcloning steps.

pAPEX-1 (SEQ. ID No:4) is a derivative of the vector pcDNA1/Amp (Invitrogen, San Diego CA) which was modified as follows to increase protein expression in mammalian cells. First, since the intron derived from the gene encoding the SV40 small-t antigen has been shown to decrease expression of upstream coding regions (Evans and Scarpulla, 1989), this intron was removed from pcDNA1/Amp by digestion with XbaI-HpaI, followed by treatment with the Klenow fragment of DNA polymerase and all four dNTPs. The resulting blunt ended 4.2 kb fragment was gel purified and self ligated to yield a closed circular plasmid. A 5'-untranslated region adenovirus/immunoglobulin hybrid intron was introduced into the plasmid by replacing a 0.5 kb NdeI-NotI fragment with the

corresponding 0.7 kb NdeI-NotI fragment from the vector pRc/CMV7SB (obtained from Dr. Joseph Goldstein, University of Texas Southwest Medical Center, Dallas, TX). Finally, the resulting CMV promoter expression cassette was shuttled as an NdeI-SfiI fragment into the vector pGEM-4Z (Promega, Madison WI) by ligation to an NdeI-SfiI fragment (containing pGEM-4Z) obtained from a pGEM based expression vector containing a CMV-promoter and an SV40 origin of replication (Davis et al., 1991).

#### Example 2

#### Transient Transfection of COS Cells with Porcine Galactose $\alpha(1,3)$ Galactosyltransferase and Human H Transferase

COS cells (ATCC # CRL 1650) were transiently transfected with CMV-based expression vectors. These vectors were pGT, containing an insert comprising a sequence (SEQ ID NO:5) encoding the pig  $\alpha(1,3)$  Gal transferase oriented for CMV promoter-driven expression in the parent vector pCDNAI (Invitrogen, San Diego, CA), and pHT (described above), encoding the human H transferase. Clones containing pig  $\alpha(1,3)$  Gal transferase cDNAs have been deposited with the Australian Government Analytical Laboratories, 1 Suakin Street, Pymble, N.S.W. 2073, Australia, and have been assigned the designations N94/9030 and N94/9029, respectively (see copending U.S. application Serial No. 08/214,580, supra). In a series of transfection experiments, the amount of pGT was kept constant at 3  $\mu$ g/well and the amount of pHT was varied between 0  $\mu$ g and 3  $\mu$ g (Table 1). Transfection was carried out using the DEAE-dextran method (Seed and Aruffo, 1987).

COS cells maintained in DMEM with 10% FBS were seeded into 6-well tissue culture plates and were subsequently transfected with pHT and/or pGT. Transfected cells were examined for the expression of the Gal  $\alpha(1,3)$  Gal epitope or the H epitope 48 hours after transfection. The cell surface expression of these two

epitopes was assessed using a fluoresceinated IB4 lectin, which binds specifically to the Gal  $\alpha(1,3)$  Gal sugar structure, or by indirect immunofluorescence using a monoclonal antibody specific for the human H epitope (ASH-1952, obtained from the Austin Research Institute, Heidelberg, Victoria, Australia; see Devine, et al., 1990) and an immunopurified, fluorescein-conjugated Sheep anti-mouse IgG (Selenus Laboratories, Melbourne, Australia) as the secondary antibody. FIGS. 1-4 show the results of phase contrast (P) and fluorescence (F) microscopy of the cells obtained in these experiments.

In particular, FIGS. 1-4 are photomicrographs of African Green Monkey COS cells which have been fluorescently stained with ASH-1952 (FIGS. 2 and 3) or with IB4 (FIGS. 1 and 4). In each figure, the bottom panel shows all cells, as seen by phase contrast illumination, and the top panel shows only those cells specifically binding to the mAb or lectin as seen by ultraviolet illumination. The cells in FIG. 1 have been transfected with pGT; the cells in FIG. 2 have been transfected with pHT; and the cells in FIGS. 3 and 4 have been transfected with equal amounts of both vectors.

To assess the percentage of cells staining for the carbohydrate epitopes, 600-800 cells from each well were counted after staining. As shown in Table 1 and in FIGS. 1-2, most COS cells transiently transfected with the porcine Gal  $\alpha(1,3)$  Gal transferase alone (3  $\mu$ g) were positive for IB4 staining, and most COS cells transfected with the human H transferase alone (3  $\mu$ g) were positive for anti-H staining. When equal amounts of the two expression plasmids were used (3  $\mu$ g each), COS cells stained predominately for the H epitope (68% of cells), with only weak staining observed for the Gal  $\alpha(1,3)$  Gal epitope (1.5% of cells). See Table 1 and FIGS. 3-4. In fact, even at a DNA ratio of 10:1 (3  $\mu$ g pGT to 0.3  $\mu$ g pHT), COS cells still stained predominately for the H

epitope (50.2 %) relative to the Gal  $\alpha(1,3)$  Gal epitope (17.5%).

As a control, cotransfections were also done using expression vectors derived from a parent CMV-based expression vector (pCDM8; Seed and Aruffo, 1987) encoding either Ly-9 (Sandrin et al., 1992) or CD48 (Vaughan et al., 1991). Staining for the Ly-9 epitope was carried out using monoclonal antibody anti-Ly-9.2 (Sandrin et al., 1992). Staining for the CD48 epitope was carried out using an anti-CD48 monoclonal antibody (HuLy-m3; Vaughan et al., 1991). When equal amounts of the Ly-9 or CD48 expression vectors were cotransfected with either pGT or pHT, COS cells demonstrated intense staining for the appropriate carbohydrate epitope and for either CD48 or Ly-9, respectively. These results indicate that two different CMV-based expression vectors can function equally well when cotransfected into COS cells.

#### Example 3

##### Stable expression of H transferase in xenogeneic cells results in down-regulation of Gal $\alpha(1,3)$ Gal expression

A porcine kidney cell line (LLC-PK<sub>1</sub>: ATCC# CRL 1392) was transfected with plasmid pHT (directing the expression of H transferase) and plasmid pSV2neo (directing the expression of the neomycin resistance gene encoding neomycin phosphotransferase) at a molar ratio of 20:1. Transfection was carried out by the calcium phosphate co-precipitation method and transfected cells were cultured in DMEM + 10% fetal bovine serum + G418 (500 $\mu$ g/ml, active). Stable neomycin resistant colonies were selected and expanded.

The cell surface expression of the H epitope was analyzed on G418 resistant colonies by indirect immunofluorescence performed with ASH-1952 (identified as "anti-H mAB" in FIG. 5) or with the H epitope specific lectin UEAI (EY Laboratories, Inc., San Mateo, CA.) directly conjugated to FITC. The Gal  $\alpha(1,3)$  Gal cell surface epitope was visualized by staining control and

transfected cells with the FITC-conjugated lectin, IB4 (EY Laboratories, Inc., San Mateo, CA). As a control, transfected LLC-PK<sub>1</sub> cells were also stained with the anti-SLA class I (anti-pig major histocompatibility antigen class I) mAb, PT85A (VMRD, Inc., Pullman WA), as a positive control. Goat anti-mouse IgG antisera (monoclonal sera, Zymed Laboratories, South San Francisco, CA) directly conjugated to FITC was used to detect specific antibody binding to the cell surface by flow cytometry as shown in FIG. 5.

These data demonstrate that G418 resistant control LLC-PK<sub>1</sub> cells (clone PK1:neo #B6) normally express low levels of both the Gal  $\alpha(1,3)$  Gal epitope and the H epitope (FIG. 5B) compared to staining with secondary antibody alone (FIG. 5D; 2° curve). However, cells transfected with the human H transferase vector (clone #A3) express high levels of the H epitope (FIG. 5A) and reduced levels of the Gal  $\alpha(1,3)$  Gal epitope (FIG. 5A). Transfection of these cells with H transferase, however, did not alter the cell surface expression of the SLA class I gene product (FIG. 5C) relative to G418 resistant control cells (FIG. 5D).

#### Example 4

##### Stable expression of H transferase in xenogeneic cells results in significantly reduced binding of human IgG and IgM antibodies

Cell surface reactivity of human serum on the LLC-PK<sub>1</sub> transfectants was measure by incubation with 0% or 20% human whole serum followed by incubation with FITC conjugated goat anti-human antibodies specific for either human IgG or human IgM (Zymed Laboratories, South San Francisco, CA). Cell surface antibody binding was then measured by flow cytometry on a FACSort instrument (Becton-Dickinson Immunocytometry Systems, San Jose, CA). As shown in FIG. 6, LLC-PK<sub>1</sub> cells stably transfected with pHT demonstrate little to no reactivity to either human IgG (FIG. 6A) or IgM (FIG. 6B) relative to G418 resistant

control cells which demonstrate significant binding to human IgG (FIG. 6C) and IgM (FIG. 6D) present in 20% human serum. The binding of human IgG and IgM present in 20% human serum to H transferase-expressing LLC-PK<sub>1</sub> cells is similar to the binding observed with 0% whole human serum. These data together with the data presented in Example 3 indicate that expression of the H epitope on the surface of the LLC-PK<sub>1</sub> cells results in down-regulation of the expression of the Gal  $\alpha(1,3)$  Gal epitope to such low levels that preformed naturally occurring human antibodies no longer bind to the cells.

#### Example 5

#### Stable expression of H transferase in xenogeneic cells results in significantly reduced sensitivity to human complement

The functional significance of recombinant H transferase expression by LLC-PK<sub>1</sub> cells was assessed by measuring the efflux of the trapped cytoplasmic indicator dye, Calcein AM (Molecular Probes, Inc.), from cells subjected to human complement-mediated damage by human serum. Transfected cells expressing the human H transferase and the neomycin resistance gene (clone #A3; see Examples 3 and 4 above) or the neomycin resistance gene alone (clone #C6; prepared in the same manner as clone B6 described above in Examples 3 and 4) were grown to confluence in 96-well plates. Cells were washed 2X with 200  $\mu$ l of HBSS containing 1% (w/v) BSA (HBSS/BSA). Calcein AM was added (10mM final) and the plates were incubated at 37°C for 30 minutes. Subsequently, the cells were incubated at 37°C for 30 minutes in the presence of increasing concentrations of human whole serum.

Dye released from the cells was determined by the fluorescence in the supernatant. Total cell associated dye was determined from a 1% SDS cell lysate. The dye release was calculated as a percent of total, correcting for non-specific dye release and background fluorescence

measured for identically matched controls without the addition of serum. Fluorescence was measured using a Millipore Cytofluor 2350 fluorescence plate reader (490nm excitation, 530nm emission).

5 As shown in FIG. 7, LLC-PK<sub>1</sub> cells stably transfected with pHT (clone #A3; open triangles) were significantly less sensitive to the lytic activity of human complement relative to control LLC-PK<sub>1</sub> cells (clone #C6; closed circles) at all concentrations of human serum tested  
10 between 1% and 40%.

Throughout this application various publications, patents, and patent applications are referred to. The teachings and disclosures of these publications, patents, and patent applications in their entireties are hereby  
15 incorporated by reference into this application to more fully describe the state of the art to which the present invention pertains.

Although preferred and other embodiments of the invention have been described herein, further embodiments  
20 may be perceived by those skilled in the art without departing from the scope of the invention as defined by the following claims.

TABLE 1

<u>pGT (<math>\mu</math>g)</u>	<u>pHT (<math>\mu</math>g)</u>	<u>IB4</u> % positive	<u>ASH-1952</u> % positive
0.0	3.0	.01	69.0
3.0	3.0	1.5	68.0
3.0	1.5	4.5	70.3
3.0	1.0	4.6	65.8
3.0	0.3	17.5	50.2
3.0	0.15	43.4	34.0
3.0	0.03	61.5	28.9
3.0	0.0	68.4	0.0

-----  
pGT = porcine galactose  $\alpha$ (1,3) galactosyltransferase cDNA subcloned into CMV-based expression plasmid pCDNAI (Invitrogen, Sand Diego, CA).

pHT = human H transferase cDNA subcloned into CMV-based expression plasmid pAPEX-1.

IB4 binds to the Gal  $\alpha$ (1,3) Gal epitope.

ASH-1952 binds to the H epitope.



REFERENCES

- Auchincloss, 1988. Transplantation 46, pp. 1.
- Ausubel et al., 1992. Current Protocols in Mol Bio, Volume 1, Section III (units 9.10.1 - 9.14.3). John Wiley & Sons, New York.
- Beyer and Hill, 1982, in Horowitz, (ed.). The Glycoconjugates Volume 3. Academic Press, NY, pp. 25-45.
- Blanken and Van den Eijnden, 1985. J Biol Chem 260, pp. 12927-12934.
- Boggs SS, Int. J. Cell Cloning, 8:80-96; 1990.
- Bradley, 1987, in Robertson (ed). Teratocarcinomas and Embryonic Stem Cells a Practical Approach. IRL Press, Eynsham, Oxford, England.
- Brent, 1991. Curr Opin Immunol 3, pp. 707.
- Brinster et al., 1985. Proc Natl Acad Sci, USA 82, pp. 4438-4442.
- Cairns et al., 1993A. Second International Congress on Xenotransplantation, pp. 56 (abstr).
- Cairns et al., 1993B. Second International Congress on Xenotransplantation, pp. 156 (abstr).
- Cone and Mulligan, Proc. Natl. Acad. Sci. USA, 81:6349-6353 1984.
- Cooper et al., 1991, in Cooper et al., (eds). Xenotransplantation Heidelberg, Springer-Verlag, pp. 481-500.
- Cosset et al., 1990. J Virol 64, pp. 1070-1078.
- Dabkowski et al., 1993. Transplant Procs 25, pp. 2921.
- Dalmaso et al., 1992. Am J Path 140, pp. 1157.
- Davis et al., 1991. Science 253, pp. 59.
- Devine, et al., 1990. J Tumor Marker Oncol 5(4), pp. 321-339.
- Eglitis et al., 1988. Biotechniques 6, pp. 608-614.
- Eglitis, 1991. Human Gene Therapy, 2:195-201.
- Evans and Scarpulla, 1989. Gene 84, pp. 135.

- Fabian et al., 1992. Transplant Procs 24, pp. 604.
- Felgner PL, Rhodes G, Nature, 349:351-352; 1991.
- Fournier et al., 1993. Second International Congress on Xenotransplantation, p 142 (abstr).
- Galili et al., 1984. J Exp Med 160, pp. 1519.
- Galili et al., 1985. J Exp Med 162, pp. 573.
- Galili et al., 1987. Proc Natl Acad Sci, USA 84, pp. 1369.
- Galili et al., 1988. J Biol Chem 263, pp. 17755-17762.
- Galili, 1993. Immunology Today 14, pp. 480.
- Geller et al., 1993. Transplantation 55, pp. 168.
- Gilboa, Biotechniques, 4:504-512, 1986.
- Good et al., 1992. Transplant Procs 24, pp. 558.
- Hammer et al., 1992. Transplant Procs 24, pp. 590.
- Hogan et al., 1986. Manipulating the Mouse Embryo: A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Kohn DB, Anderson WF, Blaese RM, Cancer Investigation, 7:179-192;1989.
- Koren et al., 1992. Transplant Proc 24, pp. 598.
- Koren et al., 1993. Second International Congress on Xenotransplantation, pp. 178 (abstr).
- Korman et al., 1987. Proc Natl Acad Sci, USA 84, pp. 2150-2154.
- Kornfeld and Kornfeld, 1985. Ann Rev Biochem 54, pp. 631-664.
- Kukowska et al., 1990. Genes Dev 4, pp. 11288-13039.
- Larsen et al., 1990a. J Biol Chem 265, pp. 7055-7061.
- Larsen et al., 1990b. Proc Natl Acad Sci USA 87, pp. 6674-6678.
- Lehn PM, Bone Marrow Transplantation, 5:287-293;1990.

- Lovell-Badge, 1987, in Robertson (ed). Teratocarcinomas and Embryonic Stem Cells a Practical Approach. IRL Press, Eynsham, Oxford, England.
- Lowe, 1991. Sem Cell Biol 2, pp. 289-307.
- Mann, et al., Cell, 33:153-159, 1983.
- Markowitz et al., 1988. J Virol 62, pp. 1120-1124.
- McMahon et al., 1990. Cell 62, pp. 1073-1085.
- Mejia-Laguna et al., 1972. Am J Path 69, pp. 71.
- Miller et al., 1986. Mol Cell Bio 6, pp. 2895-2902.
- Miller et al., 1989. Biotechniques 7, pp. 981-990.
- Miller, 1992, Nature, 357:455-460.
- Morgenstern et al., 1990. Nucleic Acids Res 18, pp. 3587-3596.
- Muler-Eberhard, 1988. Ann Rev Biochem 57, pp. 321.
- Mulligan, 1993, Science, 260:926-932.
- Najarian, 1992. Transplant Proc 24, pp. 733.
- Niekrasz et al., 1992. Transplant Procs 24, pp. 625-626.
- Pedersen et al., 1990. Transgenic Techniques in Mice - A Video Guide. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Platt and Bach, 1991. Curr Opin Immunol 3, pp. 735.
- Platt et al., 1990. Immunology Today 11, pp. 450.
- Platt et al., 1991. Transplantation 52, pp. 214.
- Robertson et al., 1986. Nature 323, pp. 445-448.
- Robertson, 1987, in Robertson (ed). Teratocarcinomas and Embryonic Stem Cells a Practical Approach. IRL Press, Eynsham, Oxford, England.
- Sambrook et al., 1989. Molecular Cloning: A Laboratory Manual, 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sandrin et al., 1992. J Immunol 149, pp. 1636-1641.
- Sandrin et al., 1993A. Proc Natl Acad Sci, USA 90, pp. 11391-11395.

- Sandrin et al., 1993B. Transplant Proc 25, pp. 2917.
- Seed and Aruffo, 1987. Proc Natl Acad Sci, USA 84, pp. 3365.
- Somerville and d'Apice, 1993. Kidney Intl 44, pp. 112.
- Stanley, 1992. Glycobiol 2(2), pp. 99-107.
- Talib et al., 1991. Gene 98, pp. 289-293.
- Thall and Galili, 1990. Biochemistry 29, pp. 3959.
- Tusso et al., 1992. Transplant Procs 24, pp. 596.
- Tusso et al., 1993. Transplantation 55, pp. 1375.
- Vaughan et al., 1991. Immunogenetics 33, pp. 113-117.
- Verma IM, Scientific American, pp. 68-84;1990.
- Weatherall DJ, Nature, 349:275-276; 1991.
- Weston et al., 1992. J Biol Chem 267, pp. 4152-4160.
- Williams et al., 1988. Immunogenetics 27, pp. 265-272.
- Yamakawa and Nagai, 1978. Trends Biol Sci 3, pp. 128-131.
- Zehr et al., 1994. Transplantation 57, pp. 900.
- Zhao and Wong, 1991. J Cell Biol 115, pp. 83a.
- Zhao et al., 1993. Second International Congress on Xenotransplantation, pp. 138 (abstr).

SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Sandrin, Mauro S.  
Fodor, William L.  
Rother, Russell P.  
Squinto, Stephen P.  
McKenzie, Ian F. C.
- (ii) TITLE OF INVENTION: Methods for Reducing  
Hyperacute Rejection of Xenografts
- (iii) NUMBER OF SEQUENCES: 5
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Maurice M. Klee
  - (B) STREET: 1951 Burr Street
  - (C) CITY: Fairfield
  - (D) STATE: Connecticut
  - (E) COUNTRY: USA
  - (F) ZIP: 06430
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: 3.5 inch, 750 Kb storage
  - (B) COMPUTER: Dell 486/50
  - (C) OPERATING SYSTEM: DOS 6.2
  - (D) SOFTWARE: WordPerfect 6.0
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 08/260,201
  - (B) FILING DATE: June 15, 1994
  - (C) CLASSIFICATION:

(D) APPLICATION NUMBER: 08/278,282

(E) FILING DATE: July 21, 1994

(F) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Klee, Maurice M.

(B) REGISTRATION NUMBER: 30,399

(C) REFERENCE/DOCKET NUMBER: ALX-144.1PCT

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (203) 255-1400

(B) TELEFAX: (203) 254-1101

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 bases
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: Oligonucleotide primer

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGCCACGAAA AGCGGACTGT GGATCCGCCA CCTG

34

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 bases
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

## (ii) MOLECULE TYPE: Other nucleic acid

## (A) DESCRIPTION: Oligonucleotide primer

## (iii) HYPOTHETICAL: No

## (iv) ANTI-SENSE: Yes

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CAGGAACACC ACCAAGCTTC TCGAGAAGAT GCCAGGCC

38



(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1174 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Double

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

(A) DESCRIPTION: Human H-transferase

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(x) PUBLICATION INFORMATION:

(A)           AUTHORS:   Larsen, R.D.  
                              Ernst, L.K.  
                              Nair, R.P.  
                              Lowe, J.B.

(B) TITLE: Molecular cloning, sequence, and expression of a human GDP-L-fucose: -D-galactoside 2-alpha-L-fucosyltransferase cDNA that can form the H blood group antigen.

(C) JOURNAL: Proceedings of the National Academy of Sciences, USA

(D) VOLUME: 87

(F) PAGES: 6674-6678

(G) DATE: SEP-1990

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

										CAAGCAGCTC	GGCC	14		
ATG	TGG	CTC	CGG	AGC	CAT	CGT	CAG	CTC	TGC	CTG	GCC	TTC	CTG	56
Met	Trp	Leu	Arg	Ser	His	Arg	Gln	Leu	Cys	Leu	Ala	Phe	Leu	
1				5					10					

CTA GTC TGT GTC CTC TCT GTA ATC TTC TTC CTC CAT ATC CAT Leu Val Cys Val Leu Ser Val Ile Phe Phe Leu His Ile His 15 20 25	98
CAA GAC AGC TTT CCA CAT GGC CTA GGC CTG TCG ATC CTG TGT Gln Asp Ser Phe Pro His Gly Leu Gly Leu Ser Ile Leu Cys 30 35 40	140
CCA GAC CGC CGC CTG GTG ACA CCC CCA GTG GCC ATC TTC TGC Pro Asp Arg Arg Leu Val Thr Pro Pro Val Ala Ile Phe Cys 45 50 55	182
CTG CCG GGT ACT GCG ATG GGC CCC AAC GCC TCC TCT TCC TGT Leu Pro Gly Thr Ala Met Gly Pro Asn Ala Ser Ser Ser Cys 60 65 70	224
CCC CAG CAC CCT GCT TCC CTC TCC GGC ACC TGG ACT GTC TAC Pro Gln His Pro Ala Ser Leu Ser Gly Thr Trp Thr Val Tyr 75 80	266
CCC AAT GGC CGG TTT GGT AAT CAG ATG GGA CAG TAT GCC ACG Pro Asn Gly Arg Phe Gly Asn Gln Met Gly Gln Tyr Ala Thr 85 90 95	308
CTG CTG GCT CTG GCC CAG CTC AAC GGC CGC CGG GCC TTT ATC Leu Leu Ala Leu Ala Gln Leu Asn Gly Arg Arg Ala Phe Ile 100 105 110	350
CTG CCT GCC ATG CAT GCC GCC CTG GCC CCG GTA TTC CGC ATC Leu Pro Ala Met His Ala Ala Leu Ala Pro Val Phe Arg Ile 115 120 125	392
ACC CTG CCC GTG CTG GCC CCA GAA GTG GAC AGC CGC ACG CCG Thr Leu Pro Val Leu Ala Pro Glu Val Asp Ser Arg Thr Pro 130 135 140	434
TGG CGG GAG CTG CAG CTT CAC GAC TGG ATG TCG GAG GAG TAC Trp Arg Glu Leu Gln Leu His Asp Trp Met Ser Glu Glu Tyr 145 150	476
GCG GAC TTG AGA GAT CCT TTC CTG AAG CTC TCT GGC TTC CCC Ala Asp Leu Arg Asp Pro Phe Leu Lys Leu Ser Gly Phe Pro 155 160 165	518
TGC TCT TGG ACT TTC TTC CAC CAT CTC CGG GAA CAG ATC CGC Cys Ser Trp Thr Phe Phe His His Leu Arg Glu Gln Ile Arg 170 175 180	560
AGA GAG TTC ACC CTG CAC GAC CAC CTT CGG GAA GAG GCG CAG Arg Glu Phe Thr Leu His Asp His Leu Arg Glu Glu Ala Gln 185 190 195	602
AGT GTG CTG GGT CAG CTC CGC CTG GGC CGC ACA GGG GAC CGC Ser Val Leu Gly Gln Leu Arg Leu Gly Arg Thr Gly Asp Arg 200 205 210	644

CCG	CGC	ACC	TTT	GTC	GGC	GTC	CAC	GTG	CGC	CGT	GGG	GAC	TAT	686
Pro	Arg	Thr	Phe	Val	Gly	Val	His	Val	Arg	Arg	Gly	Asp	Tyr	
				215					220					
CTG	CAG	GTT	ATG	CCT	CAG	CGC	TGG	AAG	GGT	GTG	GTG	GGC	GAC	728
Leu	Gln	Val	Met	Pro	Gln	Arg	Trp	Lys	Gly	Val	Val	Gly	Asp	
225					230					235				
AGC	GCC	TAC	CTC	CGG	CAG	GCC	ATG	GAC	TGG	TTC	CGG	GCA	CGG	770
Ser	Ala	Tyr	Leu	Arg	Gln	Ala	Met	Asp	Trp	Phe	Arg	Ala	Arg	
	240					245					250			
CAC	GAA	GCC	CCC	GTT	TTC	GTG	GTC	ACC	AGC	AAC	GGC	ATG	GAG	812
His	Glu	Ala	Pro	Val	Phe	Val	Val	Thr	Ser	Asn	Gly	Met	Glu	
		255					260					265		
TGG	TGT	AAA	GAA	AAC	ATC	GAC	ACC	TCC	CAG	GGC	GAT	GTG	ACG	854
Trp	Cys	Lys	Glu	Asn	Ile	Asp	Thr	Ser	Gln	Gly	Asp	Val	Thr	
			270					275					280	
TTT	GCT	GGC	GAT	GGA	CAG	GAG	GCT	ACA	CCG	TGG	AAA	GAC	TTT	896
Phe	Ala	Gly	Asp	Gly	Gln	Glu	Ala	Thr	Pro	Trp	Lys	Asp	Phe	
				285					290					
GCC	CTG	CTC	ACA	CAG	TGC	AAC	CAC	ACC	ATT	ATG	ACC	ATT	GGC	938
Ala	Leu	Leu	Thr	Gln	Cys	Asn	His	Thr	Ile	Met	Thr	Ile	Gly	
295					300					305				
ACC	TTC	GGC	TTC	TGG	GCT	GCC	TAC	CTG	GCT	GGC	GGA	GAC	ACT	980
Thr	Phe	Gly	Phe	Trp	Ala	Ala	Tyr	Leu	Ala	Gly	Gly	Asp	Thr	
	310					315					320			
GTC	TAC	CTG	GCC	AAC	TTC	ACC	CTG	CCA	GAC	TCT	GAG	TTC	CTG	1022
Val	Tyr	Leu	Ala	Asn	Phe	Thr	Leu	Pro	Asp	Ser	Glu	Phe	Leu	
		325					330					335		
AAG	ATC	TTT	AAG	CCG	GAG	GCG	GCC	TTC	CTG	CCC	GAG	TGG	GTG	1064
Lys	Ile	Phe	Lys	Pro	Glu	Ala	Ala	Phe	Leu	Pro	Glu	Trp	Val	
			340					345					350	
GGC	ATT	AAT	GCA	GAC	TTG	TCT	CCA	CTC	TGG	ACA	TTG	GCT	AAG	1106
Gly	Ile	Asn	Ala	Asp	Leu	Ser	Pro	Leu	Trp	Thr	Leu	Ala	Lys	
				355					360					
CCT	TGAGAGCCAG	GGAGACTTTC	TGAAGTAGCC	TGATCTTTCT										1149
Pro														
365														
AGAGCCAGCA	GTACGTGGCT	TCAGA												1174

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4059 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Double

(D) TOPOLOGY: Circular

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: Apex-1 Eukaryotic  
(CMV) Expression Vector

(iv) ANTI-SENSE: No

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ACGCGTTGAC ATTGATTATT GACTAGTTAT TAATAGTAAT CAATTACGGG	50
GTCATTAGTT CATAGCCCAT ATATGGAGTT CCGCGTTACA TAACTTACGG	100
TAAATGGCCC CGCCTGGCTG ACCGCCCAAC GACCCCGCC CATTGACGTC	150
AATAATGACG TATGTTCCCA TAGTAACGCC AATAGGGACT TTCCATTGAC	200
GTCAATGGGT GGA CTATTTA CGGTAAACTG CCCACTTGGC AGTACATCAA	250
GTGTATCATA TGCCAAGTAC GCCCCCTATT GACGTCAATG ACGGTAAATG	300
GCCCGCCTGG CATTATGCCC AGTACATGAC CTTATGGGAC TTTCTACTT	350
GGCAGTACAT CTACGTATTA GTCATCGCTA TTACCATGGT GATGCGGTTT	400
TGGCAGTACA TCAATGGGCG TGGATAGCGG TTTGACTCAC GGGGATTTCC	450
AAGTCTCCAC CCCATTGACG TCAATGGGAG TTTGTTTTTG CACCAAAATC	500
AACGGGACTT TCCAAAATGT CGTAACAAC TCGCCCCATT GACGCAAATG	550
GGCGGTAGGC GTGTACGGTG GGAGGTCTAT ATAAGCAGAG CTCGTTTAGT	600
GAACCGTCAG AATTCTGTTG GGCTCGCGGT TGATTACAAA CTCTTCGCGG	650
TCTTTCCAGT ACTCTTGGAT CGGAAACCCG TCGGCCTCCG AACGGTACTC	700
CGCCACCGAG GGACCTGAGC GAGTCCGCAT CGACCGGATC GGAAAACCTC	750

TCGACTGTTG	GGGTGAGTAC	TCCCTCTCAA	AAGCGGGCAT	GACTTCTGCG	800
CTAAGATTGT	CAGTTTCCAA	AAACGAGGAG	GATTTGATAT	TCACCTGGCC	850
CGCGGTGATG	CCTTTGAGGG	TGGCCGCGTC	CATCTGGTCA	GAAAAGACAA	900
TCTTTTTTGT	GTCAAGCTTG	AGGTGTGGCA	GGCTTGAGAT	CTGGCCATAC	950
ACTTGAGTGA	CAATGACATC	CACTTTGCCT	TTCTCTCCAC	AGGTGTCCAC	1000
TCCCAGGTCC	AACTGCAGGT	CGACCGGCTT	GGTACCGAGC	TCGGATCCAC	1050
TAGTAACGGC	CGCCAGTGTG	CTGGAATTCT	GCAGATATCC	ATCACACTGG	1100
CGGCCGCTCG	AGCATGCATC	TAGAAGTTGT	TTATTGCAGC	TTATAATGGT	1150
TACAAATAAA	GCAATAGCAT	CACAAATTTT	ACAAATAAAG	CATTTTTTTT	1200
ACTGCATTCT	AGTTGTGGTT	TGTCCAAACT	CATCAATGTA	TCTTATCATG	1250
TCTGGATCGA	TCCCGCCATG	GTATCAACGC	CATATTTCTA	TTTACAGTAG	1300
GGACCTCTTC	GTTGTGTAGG	TACCGCTGTA	TTCCTAGGGA	AATAGTAGAG	1350
GCACCTTGAA	CTGTCTGCAT	CAGCCATATA	GCCCCGCTG	TTCGACTTAC	1400
AAACACAGGC	ACAGTACTGA	CAAACCCATA	CACCTCCTCT	GAAATACCCA	1450
TAGTTGCTAG	GGCTGTCTCC	GAATCATTA	CACCCTCCAA	AGTCAGAGCT	1500
GTAATTTTCG	CATCAAGGGC	AGCGAGGGCT	TCTCCAGATA	AAATAGCTTC	1550
TGCCGAGAGT	CCCGTAAGGG	TAGACACTTC	AGCTAATCCC	TCGATGAGGT	1600
CTACTAGAAT	AGTCAGTGCG	GCTCCCATTT	TGAAAATTCA	CTTACTTGAT	1650
CAGCTTCAGA	AGATGGCGGA	GGGCCTCCAA	CACAGTAATT	TTCTTCCCGA	1700
CTCTTAAAT	AGAAAATGTC	AAGTCAGTTA	AGCAGGAAGT	GGACTAACTG	1750
ACGCAGCTGG	CCGTGCGACA	TCCTCTTTTA	ATTAGTTGCT	AGGCAACGCC	1800
CTCCAGAGGG	CGTGTGGTTT	TGCAAGAGGA	AGCAAAAGCC	TCTCCACCCA	1850
GGCCTAGAAT	GTTTCCACCC	AATCATTACT	ATGACAACAG	CTGTTTTTTT	1900
TAGTATTAAG	CAGAGGCCGG	GGACCCCTGG	GCCCGCTTAC	TCTGGAGAAA	1950
AAGAAGAGAG	GCATTGTAGA	GGCTTCCAGA	GGCAACTTGT	CAAAACAGGA	2000
CTGCTTCTAT	TTCTGTCACA	CTGTCTGGCC	CTGTCACAAG	GTCCAGCACC	2050
TCCATACCCC	CTTTAATAAG	CAGTTTGGGA	ACGGGTGCGG	GTCTTACTCC	2100
GCCCATCCCG	CCCCTAACTC	CGCCAGTTTC	CGCCCATTTCT	CCGCCCCATG	2150

GCTGACTAAT TTTTTTTTATT TATGCAGAGG CCGAGGCCGC CTCGGCCTCT	2200
GAGCTATTCC AGAAGTAGTG AGGAGGCTTT TTTGGAGGCC TAGGCTTTTG	2250
CAAAAAGGAG CTCCCAGCAA AAGGCCAGGA ACCGTAAAAA GGCCGCGTTG	2300
CTGGCGTTTT TCCATAGGCT CCGCCCCCCT GACGAGCATC AAAAAAATCG	2350
ACGCTCAAGT CAGAGGTGGC GAAACCCGAC AGGACTATAA AGATACCAGG	2400
CGTTTCCCCC TGGAAGCTCC CTCGTGCGCT CTCCTGTTCC GACCCTGCCG	2450
CTTACCGGAT ACCTGTCCGC CTTTCTCCCT TCGGGAAGCG TGGCGCTTTC	2500
TCAATGCTCA CGCTGTAGGT ATCTCAGTTC GGTGTAGGTC GTTCGCTCCA	2550
AGCTGGGCTG TGTGCACGAA CCCCCCGTTC AGCCCGACCG CTGCGCCTTA	2600
TCCGGTAACT ATCGTCTTGA GTCCAACCCG GTAAGACACG ACTTATCGCC	2650
ACTGGCAGCA GCCACTGGTA ACAGGATTAG CAGAGCGAGG TATGTAGGCG	2700
GTGCTACAGA GTTCTTGAAG TGGTGGCCTA ACTACGGCTA CACTAGAAGG	2750
ACAGTATTTG GTATCTGCGC TCTGCTGAAG CCAGTTACCT TCGGAAAAAG	2800
AGTTGGTAGC TCTTGATCCG GCAAACAAAC CACCGCTGGT AGCGGTGGTT	2850
TTTTTGTTTG CAAGCAGCAG ATTACGCGCA GAAAAAAGG ATCTCAAGAA	2900
GATCCTTTGA TCTTTTCTAC GGGGTCTGAC GCTCAGTGGA ACGAAAATC	2950
ACGTTAAGGG ATTTTGGTCA TGAGATTATC AAAAAGGATC TTCACCTAGA	3000
TCCTTTTAAA TTA AAAATGA AGTTTTTAAAT CAATCTAAAG TATATATGAG	3050
TAAACTTGGT CTGACAGTTA CCAATGCTTA ATCAGTGAGG CACCTATCTC	3100
AGCGATCTGT CTATTTTCGTT CATCCATAGT TGCCTGACTC CCCGTCGTGT	3150
AGATAACTAC GATACGGGAG GGCTTACCAT CTGGCCCCAG TGCTGCAATG	3200
ATACCGCGAG ACCCACGCTC ACCGGCTCCA GATTTATCAG CAATAAACCA	3250
GCCAGCCGGA AGGGCCGAGC GCAGAAGTGG TCCTGCAACT TTATCCGCCT	3300
CCATCCAGTC TATTAATTGT TGCCGGGAAG CTAGAGTAAG TAGTTCGCCA	3350
GTTAATAGTT TGCGCAACGT TGTTGCCATT GCTACAGGCA TCGTGGTGTC	3400
ACGCTCGTCG TTTGGTATGG CTTTCATTCAG CTCCGGTTCC CAACGATCAA	3450
GGCGAGTTAC ATGATCCCCC ATGTTGTGCA AAAAAGCGGT TAGCTCCTTC	3500
GGTCCTCCGA TCGTTGTCAG AAGTAAGTTG GCCGCAGTGT TATCACTCAT	3550

GGTTATGGCA	GCACTGCATA	ATTCTCTTAC	TGTCATGCCA	TCCGTAAGAT	3600
GCTTTTCTGT	GACTGGTGAG	TACTCAACCA	AGTCATTCTG	AGAATAGTGT	3650
ATGCGGCGAC	CGAGTTGCTC	TTGCCCCGGCG	TCAATACGGG	ATAATACCGC	3700
GCCACATAGC	AGAACTTTAA	AAGTGCTCAT	CATTGGAAAA	CGTTCTTCGG	3750
GGCGAAAAC	CTCAAGGATC	TTACCGCTGT	TGAGATCCAG	TTCGATGTAA	3800
CCCCTCGTG	CACCCAAC	ATCTTCAGCA	TCTTTTACTT	TCACCAGCGT	3850
TTCTGGGTGA	GCAAAAACAG	GAAGGCAAAA	TGCCGCAAAA	AAGGGAATAA	3900
GGGCGACACG	GAAATGTTGA	ATACTCATA	TCTTCCTTTT	TCAATATTAT	3950
TGAAGCATTT	ATCAGGGTTA	TTGTCTCATG	AGCGGATACA	TATTTGAATG	4000
TATTTAGAAA	AATAAACAAA	TAGGGGTTC	GCGCACATTT	CCCCGAAAAG	4050
TGCCACCTG					4059

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1423 base pairs  
 (B) TYPE: Nucleic Acid  
 (C) STRANDEDNESS: Double  
 (D) TOPOLOGY: Linear

## (ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: galactosyl transferase,  
 full coding sequence

## (iii) HYPOTHETICAL: No

## (iv) ANTI-SENSE: No

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Sus scrofa*

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

CGGGGGCCAT CCCCAGCGC ACCCAGCTTC TGCCGATCAG GAGAAAATA      49
ATG AAT GTC AAA GGA AGA GTG GTT CTG TCA ATG CTG CTT GTC      91
Met Asn Val Lys Gly Arg Val Val Leu Ser Met Leu Leu Val
                    5                      10
TCA ACT GTA ATG GTT GTG TTT TGG GAA TAC ATC AAC AGA AAC      133
Ser Thr Val Met Val Val Phe Trp Glu Tyr Ile Asn Arg Asn
    15                      20                      25
CCA GAA GTT GGC AGC AGT GCT CAG AGG GGC TGG TGG TTT CCG      175
Pro Glu Val Gly Ser Ser Ala Gln Arg Gly Trp Trp Phe Pro
    30                      35                      40
AGC TGG TTT AAC AAT GGG ACT CAC AGT TAC CAC GAA GAA GAA      217
Ser Trp Phe Asn Asn Gly Thr His Ser Tyr His Glu Glu Glu
    45                      50                      55
GAC GCT ATA GGC AAC GAA AAG GAA CAA AGA AAA GAA GAC AAC      259
Asp Ala Ile Gly Asn Glu Lys Glu Gln Arg Lys Glu Asp Asn
    60                      65                      70

```



AGA Arg	GGA Gly	GAG Glu	CTT Leu	CCG Pro	CTA Leu	GTG Val	GAC Asp	TGG Trp	TTT Phe	AAT Asn	CCT Pro	GAG Glu	AAA Lys	301
				75					80					
CGC Arg	CCA Pro	GAG Glu	GTC Val	GTG Val	ACC Thr	ATA Ile	ACC Thr	AGA Arg	TGG Trp	AAG Lys	GCT Ala	CCA Pro	GTG Val	343
85					90					95				
GTA Val	TGG Trp	GAA Glu	GGC Gly	ACT Thr	TAC Tyr	AAC Asn	AGA Arg	GCC Ala	GTC Val	TTA Leu	GAT Asp	AAT Asn	TAT Tyr	385
100						105					110			
TAT Tyr	GCC Ala	AAA Lys	CAG Gln	AAA Lys	ATT Ile	ACC Thr	GTG Val	GGC Gly	TTG Leu	ACG Thr	GTT Val	TTT Phe	GCT Ala	427
		115					120					125		
GTC Val	GGA Gly	AGA Arg	TAC Tyr	ATT Ile	GAG Glu	CAT His	TAC Tyr	TTG Leu	GAG Glu	GAG Glu	TTC Phe	TTA Leu	ATA Ile	469
			130					135					140	
TCT Ser	GCA Ala	AAT Asn	ACA Thr	TAC Tyr	TTC Phe	ATG Met	GTT Val	GGC Gly	CAC His	AAA Lys	GTC Val	ATC Ile	TTT Phe	511
				145					150					
TAC Tyr	ATC Ile	ATG Met	GTG Val	GAT Asp	GAT Asp	ATC Ile	TCC Ser	AGG Arg	ATG Met	CCT Pro	TTG Leu	ATA Ile	GAG Glu	553
155					160					165				
CTG Leu	GGT Gly	CCT Pro	CTG Leu	CGT Arg	TCC Ser	TTT Phe	AAA Lys	GTG Val	TTT Phe	GAG Glu	ATC Ile	AAG Lys	TCC Ser	595
170						175					180			
GAG Glu	AAG Lys	AGG Arg	TGG Trp	CAA Gln	GAC Asp	ATC Ile	AGC Ser	ATG Met	ATG Met	CGC Arg	ATG Met	AAG Lys	ACC Thr	637
		185					190					195		
ATC Ile	GGG Gly	GAG Glu	CAC His	ATC Ile	CTG Leu	GCC Ala	CAC His	ATC Ile	CAG Gln	CAC His	GAG Glu	GTG Val	GAC Asp	679
			200					205					210	
TTC Phe	CTC Leu	TTC Phe	TGC Cys	ATT Ile	GAC Asp	GTG Val	GAT Asp	CAG Gln	GTC Val	TTC Phe	CAA Gln	AAC Asn	AAC Asn	721
				215					220					
TTT Phe	GGG Gly	GTG Val	GAG Glu	ACC Thr	CTG Leu	GGC Gly	CAG Gln	TCG Ser	GTG Val	GCT Ala	CAG Gln	CTA Leu	CAG Gln	763
225					230					235				
GCC Ala	TGG Trp	TGG Trp	TAC Tyr	AAG Lys	GCA Ala	CAT His	CCT Pro	GAC Asp	GAG Glu	TTC Phe	ACC Thr	TAC Tyr	GAG Glu	805
240						245					250			
AGG Arg	CGG Arg	AAG Lys	GAG Glu	TCC Ser	GCA Ala	GCC Ala	TAC Tyr	ATT Ile	CCG Pro	TTT Phe	GGC Gly	CAG Gln	GGG Gly	847
		255					260					265		

GAT TTT TAT TAC CAC GCA GCC ATT TTT GGG GGA ACA CCC ACT Asp Phe Tyr Tyr His Ala Ala Ile Phe Gly Gly Thr Pro Thr 270 275 280	889
CAG GTT CTA AAC ATC ACT CAG GAG TGC TTC AAG GGA ATC CTC Gln Val Leu Asn Ile Thr Gln Glu Cys Phe Lys Gly Ile Leu 285 290	931
CAG GAC AAG GAA AAT GAC ATA GAA GCC GAG TGG CAT GAT GAA Gln Asp Lys Glu Asn Asp Ile Glu Ala Glu Trp His Asp Glu 295 300 305	973
AGC CAT CTA AAC AAG TAT TTC CTT CTC AAC AAA CCC ACT AAA Ser His Leu Asn Lys Tyr Phe Leu Leu Asn Lys Pro Thr Lys 310 315 320	1015
ATC TTA TCC CCA GAA TAC TGC TGG GAT TAT CAT ATA GGC ATG Ile Leu Ser Pro Glu Tyr Cys Trp Asp Tyr His Ile Gly Met 325 330 335	1057
TCT GTG GAT ATT AGG ATT GTC AAG ATA GCT TGG CAG AAA AAA Ser Val Asp Ile Arg Ile Val Lys Ile Ala Trp Gln Lys Lys 340 345 350	1099
GAG TAT AAT TTG GTT AGA AAT AAC ATC TGACTTTAAA Glu Tyr Asn Leu Val Arg Asn Asn Ile 355	1136
TTGTGCCAGC AGTTTTCTGA ATTTGAAAGA GTATTACTCT GGCTACTTCC	1186
TCAGAGAAGT AGCACTTAAT TTTAACTTTT AAAAAAATAC TAACAAAATA	1236
CCAACACAGT AAGTACATAT TATTCTTCCT TGCAACTTTG AGCCTTGTC	1286
AATGGGAGAA TGACTCTGTA GTAATCAGAT GTAAATTCCC AATGATTTCT	1336
TATCTGCGGA ATTCCAGCTG AGCGCCGGTC GCTACCATTA CCAGTTGGTC	1386
TGGTGTCGAC GACTCCTGGA GCCCGTCAGT ATCGGCG	1423

What is claimed is:

1. A method for reducing rejection of a xenogeneic cell following transplantation into a human or an Old World primate comprising:

(a) producing a genetically altered cell by introducing an expression vector comprising a nucleic acid sequence encoding a protein having fucosyltransferase activity into a recipient cell, the introduction of said expression vector causing a substantial reduction in the binding of naturally occurring preformed human antibodies or naturally occurring preformed Old World primate antibodies to the genetically altered cell when compared to the binding of said antibodies to the recipient cell; and

(b) transplanting said genetically altered cell or a cell derived from said cell into a human or an Old World primate.

2. The method of Claim 1 wherein the genetically altered cell is an ungulate cell.

3. The method of Claim 1 wherein the genetically altered cell is a retroviral producer cell.

4. An ungulate cell which has been genetically altered by the introduction of an expression vector comprising a nucleic acid sequence encoding a protein having fucosyltransferase activity into a recipient ungulate cell, the introduction of said expression vector causing a substantial reduction in the binding of naturally occurring preformed human antibodies or naturally occurring preformed Old World primate antibodies to said genetically altered ungulate cell when compared to the binding of said antibodies to the recipient ungulate cell.


5. An ungulate cell, tissue, or organ derived from the genetically altered ungulate cell of Claim 4.

6. A retroviral packaging cell which has been genetically altered by the introduction of an expression vector comprising a nucleic acid sequence encoding a


protein having fucosyltransferase activity into a recipient cell from which the genetically altered retroviral packaging cell is derived, the introduction of said expression vector causing a substantial reduction in the binding of naturally occurring preformed human antibodies or naturally occurring preformed Old World primate antibodies to said genetically altered retroviral packaging cell when compared to the binding of said antibodies to the recipient cell from which the genetically altered retroviral packaging cell is derived.

7. A retroviral producer cell which has been genetically altered by the introduction of an expression vector comprising a nucleic acid sequence encoding a protein having fucosyltransferase activity into a recipient cell from which the genetically altered retroviral producer cell is derived, the introduction of said expression vector causing a substantial reduction in the binding of naturally occurring preformed human antibodies or naturally occurring preformed Old World primate antibodies to said genetically altered retroviral producer cell when compared to the binding of said antibodies to the recipient cell from which the genetically altered retroviral producer cell is derived.

1/7




F




PC

*Fig 1*

2/7




F




PC

*Fig. 2*




F




PC

*Fig. 3*



F




PC


*Fig. 4*




5/7

*Fig. 5*


6/7

*Fig 6*

6/7


*Fig 6*

5/7

*Fig 5*

7/7

PKI: HUMAN SERUM DYE RELEASE ASSAY  
H-Transferase vs. NEO control

*Fig 7*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/07554

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 43/04, 63/00; A61K 31/70, 48/00; C12N 15/00

US CL : 424/93.21; 435/172.3, 320.1; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/93.21; 435/172.3, 320.1; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,283,058 (FAUSTMAN) 01 February 1994, see entire document.	1-7
Y, P	Immunological Reviews, Volume 141, issued October 1994, Gustafsson et al., "Alpha-1,3-galactosyltransferase: a target for in vivo genetic manipulation in xenotransplantation", pages 59-70, see entire document.	1-7
Y	Transplantation, Volume 56, Number 6, issued December 1993, Oriol et al., "Carbohydrate antigens of pig tissues reacting with human natural antibodies as potential targets for hyperacute vascular rejection in pig-to-man organ xenotransplantation", pages 1433-1442, see entire document.	1-7

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

09 AUGUST 1995

Date of mailing of the international search report

15 SEP 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MICHAEL NEWELL

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/07554

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Lancet, Volume 342, issued 11 September 1993, Cooper et al., "Genetically engineered pigs", pages 682-683, see entire document.	1-7
Y	Transplantation, Volume 57, Number 5, issued March 1994, Ye et al., "The pig as a potential organ donor for man", pages 694-703, see entire document.	1-7
Y	Immunology Today, Volume 14, Number 10, issued 1993, Galili, "Interaction of the natural anti-Gal antibody with alpha-galactosyl epitopes: a major obstacle for xenotransplantation in humans", pages 480-482, see entire document.	1-7
Y	Transplantation, Volume 57, Number 10, issued May 1994, Kennedy et al., "Protection of porcine aortic endothelial cells from complement-mediated cell lysis and activation by recombinant human CD59", pages 1494-1501, see entire document.	1-7
Y, P	Transplantation Proceedings, Volume 27, Number 1, issued February 1995, Strahan et al., "Pig alpha-1,3-galactosyltransferase: Sequence of a full-length cDNA clone, chromosomal localisation of the corresponding gene, and inhibition of expression in cultured pig endothelial cells", pages 245-246, see entire document.	1-7
Y, P	Transplantation Proceedings, Volume 27, Number 1, issued February 1995, McKenzie et al., "Comparative studies of the major xenoantigen GAL-alpha (1,3)gal in pigs and mice", pages 247-248, see entire document.	1-7
Y, P	Transplantation Proceedings, Volume 27, Number 1, issued February 1995, Goldberg et al., "Inhibition of the human antipig xenograft reaction with soluble oligosaccharides", pages 249-250, see entire document.	1-7
Y	Proc. Natl. Acad. Sci. USA, Volume 90, issued December 1993, Sandrin et al., "Anti-pig IgM antibodies in human serum react predominantly with Gal (alpha 1,3) Gal epitopes", pages 11391-11395, see entire document.	1-7
Y	Transplantation, Volume 57, Number 6, issued March 1994, Fukushima et al., "The role of anti-pig antibody in pig-to-baboon cardiac xenotransplantation rejection", pages 923-928, see entire document.	1-7

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/07554

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	Veterinary Immunology and Immunopathology, Volume 43, issued 1994, Sachs, "The pig as a potential xenograft donor", pages 185-191, see entire document.	1-7

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/07554

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, Medline, Embase, CAPlus

search terms: Sandrin, Fodor, Squinto, Rother, McKenzie, xenograft, xenotransplant, xenogeneic, xenoantigen, transplant, fucosyltransferase, galactosyltransferase, glycosyltransferase, ungulate, pig, porcine, hyperacute rejection, (gene, genetic, DNA, nucleic acid)(W)(therapy or delivery or transfer or treatment)

